

# Pixxaro EC®

## ARYLEX™ ACTIVE

### TECHNICAL INFORMATION SHEET

Pixxaro EC® is a new herbicide for the control of broad-leaved weeds in winter cereals (wheat, durum wheat, spelt, barley, rye and triticale, and spring cereals (wheat, durum wheat, barley and rye).

Pixxaro EC is a **non-ALS**, spring-applied, wide spectrum broad-leaved weed herbicide containing ARYLEX™ Active. Regardless of weather conditions, Pixxaro EC provides robust control of competitive weeds such as cranesbill, poppy, fumitory, fat hen, cleavers and chickweed.

#### Why use Pixxaro EC?

- Provides robust weed control irrespective of the weather or weed growth pattern – if you can travel it will work
- Wide range of following crop options
- No cultivations required before sowing most crops
- No restrictions on straw / manure use or management
- Excellent multiway compatibility including: ACCase graminicides, ALS graminicides, PGRs Fungicides and Trace elements

#### Key facts

- Product Registration Number: MAPP 17545
- Active Ingredient: 12 g/litre Arylex + 280 g/litre fluroxypyr
- Pack size: 5.0 Litres
- Formulation: Emulsifiable Concentrate (Liquid)
- Maximum Individual Dose: 0.5 L/ha from 1<sup>st</sup> February (Winter cereals)  
0.5 L/ha from 1<sup>st</sup> March (Spring cereals)
- Maximum Total Dose: 0.5 L/ha
- Application Timing: Winter cereals 1<sup>st</sup> February to 30<sup>th</sup> June GS13 to GS45  
Spring cereals 1<sup>st</sup> March to 30<sup>th</sup> June GS13 to GS45
- Water Volumes: 100-400 L/ha
- Spray Quality: Medium as defined by BCPC
- Nozzles: Flat Fan, Variable Pressure  
Flat Fan, Pre-Orifice, Air Inclusion, Airtec
- Buffer Zone: LERAP B 5m reducible buffer zone (1m dry ditches)



**Dow AgroSciences**

*Solutions for the Growing World*

Use plant protection products safely. Always read the label and product information before use. For further information including warning phrases and symbols refer to label. Dow AgroSciences Limited, Latchmore Court, Brand Street, Hitchin, Hertfordshire, SG5 1NH. Tel: +44 (0) 1462 457272. ©TM Trademark of the Dow Chemical Company ("Dow") or an affiliated company of Dow. All other brand names are trademarks of other manufacturers for which proprietary rights may exist. PIXXARO EC® contains haloxifen-methyl (Arylex™ Active) and fluroxypyr. Technical Hotline: 0800 689 8899 | UKHotline@dow.com | [uk.dowagro.com](http://uk.dowagro.com)

March 2016

## Where Does Pixxaro EC fit?

### Winter Cereals

- Tank mix with spring applied ALS graminicide
- Robust & complimentary weed control: cleavers, poppy & cranesbill
- Tank mix with T0 or T1 fungicide programme
- Robust weed control irrespective of temperature & weed growth
- Apply at T0, creates tank mix flexibility at T1
- Compatible in multiway mixes with fungicides/PGRs/trace elements/herbicides

### Spring Cereals

- Tank mix with ½ rate SU or dicamba product
- Tank mix with ACCase graminicide
- Apply mid to late tillering of the cereal crop
- Extensive BLW control offering control of the most economically damaging and competitive weeds
- Flexibility of tank mix partners including graminicides, PGRs & fungicides



### More Information:

For product label, tank mix compatibility and more information on Pixxaro EC please see:

[uk.dowagro.com/products/pixxaro-ec/](http://uk.dowagro.com/products/pixxaro-ec/)

## Weed Spectrum

Weed	0.375 L/ha + adjuvant	0.5 L/ha + adjuvant
<b>Black Bindweed</b>	S<10 cm	S<15 cm
Black Nightshade	-	S<4 lf
Charlock	T	T
<b>Chickweed</b>	S< 25 cm	S<flowering
<b>Cleavers</b>	b4 flowering	b4 flowering
Clover	-	S<4 lf
<b>Cornflower</b>	MS<12 cm	b4 flowering
<b>Cranesbill</b>	S<12 cm	S<25 cm
Docks	-	S<4 lf
<b>Fat Hen</b>	S<15 cm	S<25 cm
Forget-me-not	MS<10 cm	S<10 cm
<b>Fumitory</b>	S<20 cm	b4 flowering
Groundsel	T	MS
<b>Hemp-nettle</b>	S< 10 cm	b4 flowering
Henbit Deadnettle	S<flowering	S<flowering
<b>Knotgrass</b>	MS<10 cm	S< 8 cm
Marigold	T	T
Mayweed	T	T
Nettle, Small	S	S TBC
Orache	MS	S<4lf
Pale Persicaria	MS	S<4 lf
Pennycress	S<10 cm	S<10 cm
Pansy	T	T
<b>Poppy</b>	S<10 cm	S<10 cm
<b>Ragweed</b>	S< 5 cm	S<rosette
<b>Red Dead Nettle</b>	S<18 cm	b4 flowering
Redshank	S<3 lf	S<3 lf
Scarlet Pimpernel	MS	S<flowering
<b>Shepherds Purse</b>	S<10 cm	b4 flowering
Shepherds Needle	MS	TBC
Speedwell, Common Field	T	S<4 lf, MS>4 lf
Speedwell, Ivy Leaved	MS<2 lf	S<4 lf, MS>4 lf
Spurrey	-	S<2 lf
Wild Radish	T	T
Vol Beans	S<8 lf	b4 flowering
Vol Borage		S<2 lf
Vol Linseed		S<10 cm
Vol Oilseed Rape	T	T
Vol Peas	S<8 lf	b4 flowering
Vol Potatoes	T	MS<20 cm

Key: **bold** – label weeds, S – susceptible, MS – moderately susceptible, T – tolerant

Non-label weeds listed as an indication of the effect that would be expected to be achieved based on limited data.